



THE CIEA
NOG
mouse[®]

A SUPER IMMUNE DEFICIENT MOUSE WITH
UNPARALLELED POTENTIAL FOR ENGRAFTMENT
OF HUMAN CELLS AND TISSUES

The CIEA NOG mouse®

A SUPER IMMUNE DEFICIENT MOUSE
WITH UNPARALLELED POTENTIAL FOR
ENGRAFTMENT OF HUMAN CELLS AND
TISSUES NOD.Cg-Prkdc^{scid} Il2rg^{tm1Sug}/JicTac



Compared to other immunocompromised mouse models, the CIEA NOG mouse® has a more comprehensive and severe immunodeficient phenotype, with none of the leakiness or shortened life span displayed by some other models.

Mice engrafted with human cells and tissues allow for the study of human cells under *in vivo* conditions. The CIEA NOG mouse® represents a new generation of immune deficient mice for human cell and tissue engraftment research. This severely immune compromised mouse carries the *scid* mutation and a targeted mutation of the *Il2rg* gene on the NOD/ShiJic genetic background. The functional knock out of the IL-2R γ chain results in reduction of residual innate immunity of the NOD/ShiJic background and superior engraftment of human cells and tissues compared to any other immune deficient mouse model.

TO DISCUSS YOUR NEEDS

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

CHARACTERISTICS OF THE CIEA NOG mouse®:1,5,49,73

- Lack of functional T, B and NK cells.
- Reduced complement activity.
- Dysfunctional macrophages and dendritic cells.
- NOD SIRP α allele promotes enhanced engraftment of human hematopoietic lineages.
- Deficiencies in immune signaling, including impaired cytokine production.
- Very low lymphoma development resulting in a lifespan of more than 1.5 years, comparable to other inbred strains.
- No T or B cell leakiness observed at 7 - 10 months of age.
- Does not develop diabetes.
- The Il2rg gene is sex-linked.
- Excellent model for a variety of xenograft and human cell engraftment studies.

HOW CAN THE CIEA NOG mouse® BE USED?

The unparalleled engraftment potential of human cells and tissues in this model offers an array of new opportunities for basic and translational research in areas such as:

- Xenotransplantation of human normal and cancer cell lines and tissues
- Reconstitution of the immune system for the study of the immune component in cancer as well as efficacy and mechanism of action studies for therapeutic antibodies
- Immunity and development of the immune system
- Hematopoiesis
- Stem cell research
- Infectious disease studies
- Safety assessment of new therapeutics
- Regeneration of damaged tissues
- Development of new human cell and tissue engrafted models

WHAT OTHER BENEFITS DOES THE CIEA NOG mouse® OFFER YOU?

- Excellent availability with local production in the US and Europe.
- No upfront fees or license agreements.
- Production at the Taconic Biosciences Restricted Flora™, Excluded Flora and Defined Flora health standards permits confidence in the health of the animals used in your studies.
- Researchers purchasing the CIEA NOG mouse® may receive technical support from the inventor of the model as well as Taconic's engraftment laboratory.

TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

FIGURE 1: GROWTH CHART FOR CIEA NOG mouse® PRODUCED AT RESTRICTED FLORA™

This represents a sampling from Taconic US production colonies at the Restricted Flora™ health standard. Data was collected from 2012-2013 and represents 100 animals per sex. All animals were fed NIH-31M and housed at densities appropriate for their age. The range shown is +/- two standard deviations from the mean, however note the sample size is not large enough to necessarily represent a normal population.





FIGURE 2: GROWTH CHART FOR CIEA NOG mouse® PRODUCED AT DEFINED FLORA

This represents a sampling from Taconic US production colonies at the Defined Flora health standard. Data was collected in 2013 and represents 25 animals per sex. All animals were fed NIH-31M and housed at densities appropriate for their age. The sample size for this study is too small for calculation of a standard deviation to be meaningful.



TO DISCUSS YOUR NEEDS

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

FIGURE 3: HEMOLYTIC COMPLEMENT ACTIVITY - NOD scid AND NOG mice

Defect of hemolytic complement activity in sera of NOD scid and NOG mice.
The CIEA NOG mouse® shows defects in complement activity.





FIGURE 4: NK ACTIVITIES DEFECT - NOG mice

Defect of NK activities in spleen cells of NOG mice.
The CIEA NOG mouse® lacks NK cell activity.



TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

APPLICATIONS OF THE CIEA NOG mouse®

CANCER XENOGRAPTS - CELL LINES AND PATIENT-DERIVED XENOGRAPTS (PDX)

The CIEA NOG mouse® excels at xenografts and has been used successfully to engraft a variety of solid and blood

tumors. The CIEA NOG mouse® is a natural choice for cell lines or tissue types with poor take rates in other models and is recommended as the model of choice for use with patient-derived tumors.^{47,108,110,119,121,136}

FIGURE 5: HISTOPATHOLOGICAL EVALUATION OF ESTABLISHED TUMOR LINES IN NOG MICE




FIGURE 6: DEVELOPMENT OF PATIENT-DERIVED TUMOR LINES IN THE NOG MOUSE

Figure 5 and 6: The CIEA NOG mouse® may be used to establish patient-derived tumor models. Fresh human cancer samples were subcutaneously inoculated into the flank of NOG mice. A total of 326 tumor specimens were implanted into the mice, with *in vivo* tumor models successfully generated from 54 of the tumor samples (successfully passaged *in vivo* at least three times). Samples included tumor tissue from the original tumor site as well as metastatic sites and lymph nodes adjacent to the tumor. Histopathology of the derived *in vivo* tumor models was similar to that of the original tumor sample. Data from reference 47 and IWHM 2006 meeting (Poster 18): Chen YU et al., Tokyo, Japan, Oct 11-12, 2006.

CANCER ORIGIN	NUMBER OF IMPLANTED SAMPLES			NUMBER OF SUCCESSFUL PDX MODELS GENERATED		
	PRIMARY	METASTATIC	LYMPH NODE	PRIMARY	METASTATIC	LYMPH NODE
EPITHELIAL TISSUES						
Large intestine	48	0	14	17	0	4
Stomach	18	3	2	3	1	1
Ampulla of Vater	4	0	0	2	0	0
Uterus	10	0	1	2	0	0
Ovary	18	2	0	1	1	0
Mammary gland	57	2	13	3	0	0
Cervix	4	0	2	0	0	1
Prostate	12	0	0	0	0	0
Testis	3	0	1	0	0	0
Kidney	18	0	0	3	0	0
Bladder	7	0	0	2	0	0
Lung	2	1	3	1	0	0
Pancreas	6	0	1	1	0	1
Skin	10	0	1	0	0	0
Thyroid	7	0	1	0	0	0
Others	4	1	3	0	0	0
Primary unknown	0	7	9	0	3	1
MESENCHYMAL TISSUES						
Brain	6	0	0	1	0	0
Hematopoietic	6	0	0	0	0	0
Bone	3	0	0	0	0	0
Others	16	0	0	5	0	0
TOTALS	259	16	51	41	5	8

TO DISCUSS YOUR NEEDS

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

As a highly immunodeficient model, the NOG mouse provides the best opportunity for engraftment of primary tumor samples and exerts less selection pressure which may alter tumor composition in less immunodeficient hosts. Engraftment of primary blood cancers is a key application facilitated by the NOD strain-specific Sirpa allele.⁴⁵ Male NOG mice are generally non-aggressive and can be group-housed. Thus, either male or female NOG mice may be used for general xenograft studies.

EXAMPLES OF COMPLEX CANCER MODELS GENERATED USING THE CIEA NOG mouse®

Primary adult **T-cell leukemia-lymphoma (ATL)** cells from human patients engraft successfully in the CIEA NOG mouse® via both intraperitoneal or subcutaneous implantation, resulting in infiltration by cancer cells into various organs.²⁶

Implantation of either human T-cell leukemia virus type-1 (HTLV-1)-infected cells or primary human ATL cells into the NOG results in infiltration of tumor cells

into various organs in a leukemic manner within a rapid 2-3 week period.^{11,26} In another report, investigators generated a HTLV-1 infected humanized NOG mouse by intra-bone marrow injection of human CD133+ stem cells.¹³⁸

Implantation of the U266 cell line results in a reproducible **multiple myeloma** model in the CIEA NOG mouse®, but not in C.B-17 scid or NOD scid mice. This myeloma model requires only intravenous inoculation of myeloma cells into irradiated or non-irradiated adult mice,

FIGURE 7: THE NOG MOUSE HAS BEEN SHOWN TO SUCCESSFULLY ENGRAFT MANY CANCER CELL LINES

CANCER TYPE	CELL TYPE
Breast Cancer	MCF-7 ²⁵ , MDA-MB-231 ^{25,141, 144}
Esophageal cancer	KE4 ¹¹⁶
Cholangiocarcinoma	RBE, HuCCT1 ¹⁴³
Cervical cancer	HeLa ⁶⁶
Colon cancer	COLO320DM, HCT116, HT29, LoVo, LS174T, WiDr ⁴⁸
HTLV-1 transformed T cell line	Hut-102, MT-2, MT-4, SLB-1 ¹¹
Leukemia	ED-40515(-) ¹¹ , MT-1 ¹¹ , NB4 ¹⁶ , TL-oml ¹¹ , TRL-01 ³⁴ , OSU-CLL ¹³⁷
Lung cancer	QG56 ¹¹⁶
Lymphoma	BCBL-1 ²⁷ , HDLM2 ¹⁹ , KM-H2 ¹⁹ , L428 ¹⁹ , L540 ¹⁹ , LM-2-JCK ⁵⁶ , TY-1 ²⁷ , SNK6 ¹²⁷
Melanoma	A375, A2058, G361, HMY-1 ¹⁸
Myeloma	KMM-1 ¹⁴ , U266 ^{13,14} , RPMI8226 ¹²⁸ , NCI H929 ¹²⁸
Pancreatic cancer	AsPC-1, BxPC-3, Capan-1, Capan-2, MIA PaCa-2, PL45, Panc-1 ³⁸
Pleural mesothelioma	ACC-MESO-1 ⁹⁷

FIGURE 8: THE NOG MOUSE ENGRAFTS CELL LINES OTHER MODELS CANNOT

CELL LINE	CANCER CELL TYPE	COMPARISON LINE	COMPARISON LINE TAKE RATE	NOG TAKE RATE
U266 ^{13,14}	human multiple myeloma	C.B-17 scid	0/5	20/20
U266 ^{13,14}	human multiple myeloma	NOD scid	0/5	20/20
A2058 ^{18*}	human melanoma	NOD scid	0/6	6/6
A375 ^{18*}	human melanoma	NOD scid	0/9	8/9
G361 ^{18*}	human melanoma	NOD scid	0/6	2/6
HMY-1 ^{18*}	human melanoma	NOD scid	0/8	2/8
AsPC-1 ^{38**}	human pancreatic cancer	NOD scid	8/9	9/9
BxPC-3 ^{38**}	human pancreatic cancer	NOD scid	0/8	8/8
Capan-1 ^{38**}	human pancreatic cancer	NOD scid	0/10	9/10
MIA PaCa-2 ^{38**}	human pancreatic cancer	NOD scid	0/10	10/10
PANC-1 ^{38**}	human pancreatic cancer	NOD scid	0/10	8/8

* Presence of distant metastases after iv inoculation.

** Measurement of liver metastases after intrasplenic injection.

TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

and not more difficult methods such as human fetal bone implantation or intra-osseus injection.¹³

Dissociated human **glioblastoma** cells can be implanted orthotopically in NOG mice to develop patient-specific models that recapitulate various properties of the human tumors. *In vivo* tumorigenicity correlated with clinical aggressiveness. Similarly, invasiveness, proliferation index and microvessel density of the xenograft and parental tumors were correlated. Treatment of the mouse avatars with whole brain radiation showed increased survival compared to untreated controls, and this increased survival was correlated to the progression free survival in the radiation-treated patients.¹²¹

The CIEA NOG mouse® also serves as a good model for the study of cancer **metastasis**. Four different **melanoma** cell lines, A2058, A375, G361 and HMY-1, all showed metastasis by 6 weeks post-iv injection in the CIEA NOG mouse®, whereas no metastasis was

observed in NOD scid mice.¹⁸ Metastatic potential of various **pancreatic cancer** cell lines have been assessed in the CIEA NOG mouse®,^{38,139} with significantly more metastases seen compared to the NOD scid model.³⁸ The CIEA NOG mouse® also serves as a reliable liver metastasis model for **colon cancer**.⁴⁸

In comparison to other **Hodgkin's disease** models, which either display very low engraftment rates or require conditioning such as total body irradiation or innate immunity-blocking treatments, unconditioned NOG mice efficiently engraft B-cell type Hodgkin's lymphoma cells. Tumors form within 30-60 days and retain original tumor morphology and marker expression. Depending on the route of administration, tumor cells infiltrate into regional lymph nodes or organs such as liver, lung, spleen or bone marrow.¹⁹

The superior engraftment ability of the CIEA NOG mouse® was demonstrated by dilution experiments. Indeed, as

few as 10 implanted HeLa cells formed tumors with a 50% take rate.⁶⁶ The higher susceptibility of NOG mice to xenotransplanted tumor cells indicates this model may be superior for tumorigenicity testing conducted as part of the safety assessment of stem cell therapies and other cell transplantation treatments.^{51,72}

CANCER IMMUNOTHERAPY

The CIEA NOG mouse® is an ideal choice for cancer immunotherapy experiments.^{65,79,93,116,125,129} Combine tumor engraftment with human immune system engraftment or engraftment of specific human cell types such as NK or CTL cells. Various immunotherapy approaches, including passive immunotherapy (antibody drugs and ADCC effects), active immunotherapy (cancer vaccines), and other immunomodulatory therapies can be studied using the NOG mouse.

XENOTRANSPLANTATION OF NORMAL HUMAN TISSUE

The superior engraftment potential of the CIEA NOG mouse® permits transplantation and development of normal human tissue in this model. This capacity makes the CIEA NOG mouse® a promising model for use in regenerative medicine studies. For example, the CIEA NOG mouse® can serve as a model of normal human endometrium *in vivo*, offering a more relevant model system to study human menstruation and reproduction. Ovariectomized NOG mice implanted with primary human endometrial tissue and treated with sex hormones show high levels of engraftment. This model system maintains normal tissue structure and serves as a model of the human menstrual cycle.²² Functional endometrial tissue can also be reconstituted in the CIEA NOG mouse® via implantation of singly dispersed human endometrial cells.⁴¹

FIGURE 9: THE CIEA NOG MOUSE® PROVIDES A TOOL TO EVALUATE METASTATIC CAPACITY OF DIFFERENT CELL LINES

Compared to the NOD scid model, the CIEA NOG mouse® had a far higher incidence of liver metastasis after intrasplenic injection of various pancreatic cancer cell lines. Liver metastases were dose dependent.³⁸

CELL LINE	CELL DOSE (cells/mouse)	NUMBER OF ANIMAL WITH LIVER METASTASIS Metastasis/Total	
		NOD scid	NOG
MIA PaCa-2	1x10 ⁴	0/10 (0.0%)	10/10 (100.0%)
	1x10 ³	0/7 (0.0%)	5/6 (83.3%)
	1x10 ²	0/6 (0.0%)	5/7 (71.4%)
AsPC-1	1x10 ⁴	8/9 (88.9%)	9/9 (100.0%)
	1x10 ³	2/8 (25.0%)	8/8 (100.0%)
	1x10 ²	0/6 (0.0%)	4/7 (57.1%)
PANC-1	1x10 ⁴	0/10 (0.0%)	8/8 (100.0%)
	1x10 ³	0/6 (0.0%)	6/8 (75.0%)
	1x10 ²	0/7 (0.0%)	3/8 (37.5%)
Capan-1	1x10 ⁴	0/10 (0.0%)	9/10 (90.0%)
	1x10 ³	0/10 (0.0%)	5/10 (50.0%)
	1x10 ²	0/8 (0.0%)	0/8 (0.0%)
BxPC-3	1x10 ⁵	0/8 (0.0%)	8/8 (100.0%)
	1x10 ⁴	0/8 (0.0%)	1/8 (12.5%)
Capan-2	1x10 ⁵	0/8 (0.0%)	0/8 (0.0%)
	1x10 ⁴	ND	0/10 (0.0%)
PL45	1x10 ⁵	0/8 (0.0%)	0/8 (0.0%)
	1x10 ⁴	ND	0/10 (0.0%)

HUMAN IMMUNE SYSTEM ENGRAFTMENT

As a super immune deficient model and a better recipient for engraftment of human cells, the CIEA NOG mouse® is the ideal model for human immune system engraftment experiments. The CIEA NOG mouse® may be engrafted with human hematopoietic stem cells (HSCs), which differentiate *in vivo* to form a functioning human immune system in a mouse. The NOG mouse readily accepts various sources of HSCs with high engraftment rates and successful differentiation into the various cells types of the immune system. Multiple HSC sources have been shown to engraft well in the CIEA NOG mouse®.^{5,10} In comparison to NOD/Shi-*Prkdc*^{scid} (NOD scid) mice, NOD/Shi-*Prkdc*^{scid} mice treated with anti-asialo GM1 antibody and NOD.Cg-*Prkdc*^{scid} *B2m*^{tm1Unc}/J mice, the CIEA NOG mouse® displayed higher engraftment in peripheral circulation, spleen and bone marrow compared to all other models tested.⁵

Complex experimental systems such as human/murine hybrid fetal thymic organ culture (FTOC) are not necessary for engraftment in the CIEA NOG mouse®. HSCs can simply be injected intravenously into neonatal, weanling or adult mice, with or without prior irradiation of the host. In contrast to other models, HSCs migrate with high efficiency into the thymus and peripheral lymphoid organs of the CIEA NOG mouse®.⁶

CELL TYPE DIFFERENTIATION

Nearly all cell types of the human immune system will differentiate and develop upon engraftment of the NOG mouse with human HSCs, however the interaction between those cell types may be incomplete in some cases. Production of functionally mature human T cells is possible in the CIEA NOG mouse® after implantation of HSCs, with human T cells developing in the mouse thymus, migrating to the periphery and maturing, and demonstrating production of human cytokines as well as cytotoxic capability. These differentiated T cells display a diverse V β repertoire.

Immature B cells are found primarily in bone marrow, with mature B cells in the spleen. Natural killer cells are found in both bone marrow and spleen. Human IgM, IgA and IgG from mature B cells are detectable in the sera of humanized CIEA NOG mice. In comparison, engrafted NOD.Cg-*Prkdc*^{scid} *B2m*^{tm1Unc}/J mice had little to no expression of human IgM and no expression of human IgA and IgG.⁹ Human mast cells of both the connective tissue- and mucosal-types also form in NOG mice after reconstitution with HSCs, with distribution similar to that in humans: in skin, lung, gastric tract, lymph nodes, spleen, and the peritoneal cavity.¹⁵ NOG mice reconstituted with HSCs can maintain human hematopoiesis for more than one year.⁴⁰

The BLT model, which involves surgical engraftment of autologous human liver, thymus and enriched CD34+ HSCs is a

more complex model to generate, but may result in more functional and mature human cell types. Graft versus host disease (GvHD) has not been found in the CIEA NOG mouse® after reconstitution using cord blood hematopoietic stem cells, however, implantation of human peripheral blood mononuclear cells (PBMCs) can induce graft versus host disease reliably.^{7,9,69}

Induce GvHD via engraftment of PBMCs in order to develop models of diseases involving dysfunctional T cells.¹³³ Or engraft PBMCs from patients with autoimmune diseases like lupus or Stevens-Johnson Syndrome to develop small animal models of that disease.¹¹⁴

Human cell products may be produced in human immune system engrafted NOG mice. The CIEA NOG mouse® may be reconstituted with HSCs and immunized with the peptide of interest. Hybridoma lines derived from spleen cells of immunized mice can produce a human-derived antibody against a given peptide.³⁰

Reconstitution of the CIEA NOG mouse® via cell or tissue engraftment will be an important tool for infectious disease studies as well as efficacy and safety testing of biologics and vaccines. The CIEA NOG mouse® could serve as an appropriate test species for any biologic directed at a target expressed on a human cell that can be grown in or differentiated in the CIEA NOG mouse®. This may permit the reduction of primate studies in some instances.³³

TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

MYELOABLATION

Myeloablation improves engraftment of human hematopoietic stem cells by providing a physical niche for engraftment. Myeloablation is recommended for human HSC engraftment experiments, but is not required for human PBMC engraftment. Irradiation is the most common myeloablative treatment. Use care in selecting radiation doses, as the NOG mouse (like all scid mice) is sensitive to radiation. Radiation sensitivity will vary by age and weight of the mouse. Taconic recommends users perform an irradiation dose study starting below 1 Gy to determine the most appropriate irradiation dose for your experimental setup.

Alternatives to irradiation include treatment with myeloablative drugs such as busulfan.



IMMUNE SYSTEM ENGRAFTMENT MODELS


MODEL	ENGRAFTMENT PROCEDURE	BENEFITS	LIMITATIONS	NOTES
Hematopoietic stem cells (HSCs)	Intravenous injection of CD34+ HSCs.	Good model of T cell functionality. No GvHD observed. Simple technical procedure (iv injection).	Engraftment and differentiation requires minimum of 12 weeks. Some cell types missing or in low numbers. Some cell types are not completely mature or functional.	Adults, neonates and juveniles have been used. Engraftment efficiency and function are better in neonates and juveniles than in adults, and females engraft better than males. Juvenile mice are easier to source in large, consistent cohorts, so 3-4 week old females are recommended for this application.
BLT surgical model	Co-implantation of fetal liver and thymus tissue fragments under renal capsule, plus intravenous injection of CD34+ HSCs.	Human thymus provides human thymic micro-environment for maturation of T cells. Human liver provides important human cytokines for cell maturation. Better B cell function than in HSC model.	Engraftment and differentiation requires minimum of 12 weeks. Some cell types missing or in low numbers. Some cell types are not completely mature or functional. Some incidence of GvHD possible. Complex surgical procedure required.	Use of 4 week old females is recommended.
Peripheral mononuclear blood cells (PBMCs) or purified cell subsets	Intravenous injection of peripheral blood mononuclear cells.	Ready to use quickly. Mature and functional human cells present. Simple technical procedure (iv or other type of injection).	Usage limited to -4-6 weeks only as GvHD will develop after engraftment of mature human T cells.	Male or female adult mice may be used.

TO DISCUSS YOUR NEEDS

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

FIGURE 10 FLOW CYTOMETRY ANALYSIS OF HUMAN CELL LINEAGES AFTER CD34+ HSC ENGRAFTMENT IN THE NOG MOUSE




Flow cytometry analysis of human cell lineages after CD34+ HSC engraftment.

(A) Representative flow cytometry plot of the human (hCD45+) cell to mouse cell (mCD45+) ratio at 12 weeks post-engraftment. (B) Relative percentage of human cells expressing the B cell marker CD20 vs. the T cell marker CD3. (C) Consistent engraftment ratios of greater than 25% hCD45+ cells in over 80 percent of animals (n=91), with highly reproducible relative T cell and B cell enrichment levels. Each dot represents an individual mouse analyzed by flow cytometry 12 weeks post CD34+ HSC engraftment.

FIGURE 11: A SUMMARY OF HUMAN HEMATOPOIETIC CELLS DIFFERENTIATED FROM CD34+ CELLS IN NOG MICE

A summary of human hematopoietic cell types differentiated from CD34+ cord blood cells after implantation into the CIEA NOG mouse®.



++ means dominant and well-developed; + means developed; ± means no or very few cells detected.

TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

INFECTIOUS DISEASE MODELS

HIV

The CIEA NOG mouse® serves as a unique small animal HIV model. With ongoing hematopoiesis, this model permits the study of the infectious process, as well as the study of persistent infection. When implanted with peripheral blood mononuclear cells or cord blood HSCs and infected with the R5 or X4 HIV-1 isolates, the CIEA NOG mouse® experiences a massive and systemic HIV-1 infection. In this experimental system, the mouse develops a stable and systemic infection with high viral loads and production of HIV-specific antibodies.^{20,36}

Human immune system engrafted NOG mice can be used to study HIV transmission, prophylaxis, and treatment.

EPSTEIN-BARR VIRUS

NOG mice engrafted with cord blood HSCs and infected with Epstein-Barr virus (EBV) serve as a comprehensive small animal model of human EBV infection. Infection of NOG mice engrafted with human HSCs with large doses of EBV resulted in a mostly fatal lymphoproliferative disorder, similar to

that seen in immunocompromised human patients. Lower doses of EBV induced a persistent, asymptomatic infection. The infected NOG mice developed an EBV-directed CD8+ T cell response and produced antibodies directed at EBV.⁶² In another study, humanized NOG mice infected with EBV developed erosive arthritis.¹⁰² An animal model of chronic active EBV infection was developed through injection of peripheral blood mononuclear cells isolated from patients with chronic EBV into NOG mice.¹³⁵



TO DISCUSS YOUR NEEDS

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

ORIGINS OF THE CIEA NOG mouse®

The CIEA NOG mouse® was developed by Mamoru Ito of the Central Institute for Experimental Animals (CIEA) in Japan in 2000. The *Prkdc^{scid}* mutation was identified by Mel Bosma of the Fox Chase Cancer Center in a C.B-17 congenic mouse population. This mutation was backcrossed onto the NOD/ShiJic strain at CIEA for at least eight generations. The *Il2rg* targeted mutation was developed by Kazuo Sagamura of Tohoku University by targeting the gene in ES cells derived from a 129 strain. Targeted ES cells were injected into C57BL/6 blastocysts. Resultant chimeras were backcrossed onto the C57BL/6JJic background for at least eight generations. The NOG mouse was developed by back crossing the C57BL/6JJic-*Il2rg^{tm1Sug}* line to the NOD/ShiJic-*Prkdc^{scid}* line for a total of eight generations. Taconic received stock in 2006, and the line was derived through embryo transfer. The mice are maintained by breeding females homozygous for both the *Prkdc^{scid}* and *Il2rg* mutant alleles with males that are homozygous for the *Prkdc^{scid}* allele and hemizygous for the *Il2rg* mutant allele.

THE CIEA NOG mouse® PATENTS AND TRADEMARKS

The CIEA NOG mouse® is produced and distributed under license rights to the following patents and trademarks:

- Japanese Patent No. 3,753,321
- US Patent No. 7,145,055; 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,789,215; 6,204,061; 6,653,113; 6,689,610
- EP Patent No. 1,338,198
- Japanese Trademark Reg. No. 4,823,423
- US Trademark Reg. No. 3,118,040
- EU Trademark Reg. No. 3,736,758

TRY THE CIEA NOG mouse®

Purchase NOG mice

NOG mice are produced in the US and Europe. Call Taconic Customer Service today to place an order for immediate delivery. There are no upfront fees or agreements to sign.

Purchase human immune system engrafted NOG mice

Receive mice already engrafted with a human immune system for use in further experiments in your laboratory. Skip over validation steps and go right to your experimental question. Taconic offers NOG mice engrafted with human PBMCs, human HSCs or engrafted using the BLT procedure. For more information, visit

[Taconic.com/immune-system-engrafted-mice](https://www.taconic.com/immune-system-engrafted-mice)



TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

REFERENCES DESCRIBING THE CIEA NOG mouse®

1. Ohbo K, Suda T, Hashiyama M, Mantani A, Ikebe M, Miyakawa K, Moriyama M, Nakamura M, Katsuki M, Takahashi K, Yamamura K, Sugamura K. (1996) Modulation of hematopoiesis in mice with a truncated mutant of the interleukin-2 receptor gamma chain. *Blood* 87(3):956-967.
2. Ueyama Y and Ito M. (2001) Advance in the hu-mouse. (in Japanese) *Tissue Culture Engineering* 27(7):147-151.
3. Hiramatsu H, Nakahata T. (2001) Human hematopoiesis and reconstitution of immune system in NOD/SCID mice. (in Japanese) *Cellular Molecular Medicine* 2(5):152-158.
4. Ito M. (2001) Animal model for HIV-1 infection. (in Japanese) *Sogo Rinsho* 50(10):190-192.
5. Ito M, Hiramatsu H, Kobayashi K, Suzue K, Kawahata M, Hioki K, Ueyama Y, Koyanagi Y, Sugamura K, Tsuji K, Heike T, Nakahata T. (2002) NOD/SCID/gammacnull mouse: an excellent recipient mouse model for engraftment of human cells. *Blood*, 100(9):3175-3182.
6. Saito Y, Kametani Y, Hozumi K, Mochida N, Ando K, Ito M, Nomura T, Tokuda Y, Mukuuchi H, Tajima T, Habu S. (2002) The *in vivo* development of human T cells from CD34+ cells in the murine thymic environment. *Int. Immunol.*, 14(10):1113-1124.
7. Yahata T, Ando K, Nakamura Y, Uevama Y, Shimamura K, Tamaoki N, Kato S, Hotta T. (2002) Functional human T lymphocyte development from cord blood CD34+ cells in nonobese diabetic/shi-scid, IL-2 receptor J. *Immunol.*, 169:204-209.
8. Hiramaki H, Hakahata T. (2002) Analysis of human hematopoietic stem cells using NOD/SCID/gc-/- mice. (in Japanese) *Annual Review Blood*, 159-165.
9. Hiramatsu H, Nishikomoro R, Heike T, Ito M, Kobayashi K, Katamura K, Nakahata T. (2003) Complete reconstitution of human lymphocytes from cord blood CD34+ cells using the NOD/SCID/gammacnull mice model. *Blood*, 102:873-880.
10. Matsumura T, Kametani Y, Ando K, Hirano Y, Katano I, Ito R, Shiina M, Tsukamoto H, Saito Y, Tokuda Y, Kato S, Ito M, Motoyoshi K, Habu S. (2003) Functional CD5+ B cells develop predominantly in the spleen of NOD/SCID/gammacnull (NOG) mice transplanted either with human umbilical cord blood, bone marrow and mobilized peripheral blood CD34+ cells. *Exp. Hematol.* 31:789-797.
11. Dewan Z, Terashima K, Taruishi M, Hasegawa H, Ito M, Tanaka Y, Mori N, Sata T, Koyanagi Y, Maeda M, Kubuki Y, Okayama A, Fujii M, Yamamoto N. (2003) Rapid tumor formation of human T-cell leukemia virus type 1-infected cell lines in novel NOD-SCID/gammacnull mice: suppression by an inhibitor against NF- B. *J Virol.*, 77(9):5286-5294.
12. Yahata T, Ando K, Miyatake H, Uno T, Sato T, Ito M, Kato S, Hotta T. (2004) Competitive repopulation assay of two gene-marked cord blood units in NOD/SCID/gammacnull mice. *Molecular Therapy*, 10(5).
13. Miyakawa Y, Ohnishi Y, Tomisawa M, Monnai M, Kohmura K, Ueyama Y, Ito M, Ikeda Y, Kizaki M, Nakamura M. (2004) Establishment of a new model of human multiple myeloma using NOD/SCID/gammacnull (NOG) mice. *Biochem Biophys Res Commun.*, 313:258-262.
14. Dewan Z, Watanabe M, Terashima K, Aoki M, Sata T, Honda M, Ito M, Yamaoka S, Watanabe T, Horie R, Yamamoto N. (2004) Prompt tumor formation and maintenance of constitutive NF- B activity of multiple myeloma cells in NOD/SCID/gammacnull mice. *Cancer Sci.*, 95(7):564-568
15. Kambe N, Hiramatsu H, Shimonaka M, Fujino H, Nishikomori R, Heike T, Ito M, Kobayashi K, Ueyama Y, Matsuyoshi N, Miyachi Y, Nakahata T. (2004) Development of both human connective tissue-type and mucosal-type mast cells in mice from hematopoietic stem cells with identical distribution pattern to human body. *Blood*, 103(3):860-867.
16. Ninomiya M, Kiyoi H, Ito M, Hirose Y, Naoe T. (2004) Retinoic acid syndrome in NOD/SCID mice induced by injecting an acute promyelocytic leukemia cell line. *Leukemia.*, 18:442-448.
17. Andreassen J, Ito A, Ito M, Nakao M, Nakaya K. (2004) *Hymenolepis microstoma*: direct life cycle in immunodeficient mice. *J Helminthol*, 78(1):1-5.

TO DISCUSS YOUR NEEDS

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

18. Ikoma N, Yamazaki H, Abe Y, Oida Y, Ohnishi Y, Suemizu H, Matsumoto H, Matsuyama T, Ohta Y, Ozawa A, Ueyama Y, Nakamura M. (2005) S100A4 expression with reduced E-cadherin expression predicts distant metastasis of human malignant melanoma cell lines in the NOD/SCID/gammacnull (NOG) mouse model. *Oncol Rep.*, 14:633-637.
19. Dewan MZ, Watanabe M, Ahmed S, Terashima K, Horiuchi S, Sata T, Honda M, Ito M, Watanabe T, Horie R, Yamamoto N. (2005) Hodgkin's lymphoma cells are efficiently engrafted and tumor marker CD30 is expressed with constitutive nuclear factor- κ B activity in unconditioned NOD/SCID/gammacnull *Cancer Sci.*, 96(8):466-473.
20. Nakata H, Maeda K, Miyakawa T, Shibuyama S, Matsuo M, Takaoka Y, Ito M, Koyanagi Y, Mitsuya H. (2005) Potent anti-R5 human immunodeficiency virus type 1 effects of a CCR5 antagonist, AK602/ONO4128/GW873140, in a novel human peripheral blood mononuclear cell nonobese diabetic-SCID, interleukin-2 receptor. *J. Virol.*, 79(4):2087-2096.
21. Dewan MZ, Terunuma H, Ahmed S, Ohba K, Takada M, Tanaka Y, Toi M, Yamamoto N. (2005) Natural killer cells in breast cancer cell growth and metastasis in SCID mice. *Biomed Pharmacother*, 59 Suppl 2:S375-379.
22. Matsuura-Sawada R, Murakami T, Ozawa Y, Nabeshima H, Akahira J, Sato Y, Koyanagi Y, Ito M, Terada Y, Okamura K. (2005) Reproduction of menstrual changes in transplanted human endometrial tissue in immunodeficient mice. *Hum Reprod.*, 20:1477-1484.
23. Dewan MZ, Terashima K, Ahmed S, Ohba K, Taruishi M, Yamamoto N. (2005) Mouse serum factor(s) down-modulate the CD4 and CXCR4 molecules on human T cells conferring resistance to HIV infection in NOG mice. *Med Microbiol Immunol*, 194(4):175-80.
24. Watanabe M, Dewan MZ, Okamura T, Sasaki M, Itoh K, Higashihara M, Mizoguchi H, Honda M, Sata T, Watanabe T, Yamamoto N, Umezawa K, Horie R. (2005) A novel NF- κ B inhibitor DHMEQ selectively targets constitutive NF- κ B activity and induces apoptosis of multiple myeloma cells *in vitro* and *in vivo*. *Int J Cancer*, 114(1):32-8.
25. Dewan MZ, Ahmed S, Iwasaki Y, Ohba K, Toi M, Yamamoto N. (2006) Stromal cell-derived factor-1 and CXCR4 receptor interaction in tumor growth and metastasis of breast cancer. *Biomed Pharmacother* 60:273-276.
26. Dewan MZ, Uchihara J, Terashima K, Honda M, Sata T, Ito M, Fujii N, Uozumi K, Tsukasaki K, Tomonaga M, Kubuki Y, Okayama A, Toi M, Mori N, Yamamoto N. (2006) Efficient intervention of growth and infiltration of primary adult T-cell leukemia cell by an HIV protease inhibitor, ritonavir. *Blood*, 107(2):716-724.
27. Dewan MZ, Terunuma H, Toi M, Tanaka Y, Katano H, Deng X, Abe H, Nakasone T, Mori N, Sata T, Yamamoto N. (2006) Potential role of natural killer cells in controlling growth and infiltration of AIDS-associated primary effusion lymphoma cells. *Cancer Sci.*, 97:1381-1387.
28. Dewan MZ, Terunuma H, Takada M, Tanaka Y, Abe H, Sata T, Toi M, Yamamoto N. (2006) Role of natural killer cells in hormone-independent rapid tumor formation and spontaneous metastasis of breast cancer cells *in vivo*. *Breast Cancer Res Treat.*, 104(3):267-75.
29. Doshi M, Koyanagi M, Nakahara M, Saeki K, Saeki K, Yuo A. (2006) Identification of human neutrophils during experimentally induced inflammation in mice with transplanted CD34+ cells from human umbilical cord blood. *Int J Hematol.*, 84:231-237.
30. Kametani Y, Shiina M, Katano I, Ito R, Ando K, Toyama K, Tsukamoto H, Matsumura T, Saito Y, Ishikawa D, Taki T, Ito M, Imai K, Tokuda Y, Kato S, Tamaoki N, Habu S. (2006) Development of human-human hybridoma from anti-Her-2peptide-producing B cells in immunized NOG mouse. *Exp Hematol*, 34:1239-1247.
31. Muguruma Y, Yahata T, Miyatake H, Sato T, Uno T, Itoh J, Kato S, Ito M, Hotta T, Ando K. (2006) Reconstitution of the functional human hematopoietic microenvironment derived from human mesenchymal stem cells in the murine bone marrow compartment. *Blood.*, 107:1878-1887.
32. Miyazato P, Yasunaga J, Taniguchi Y, Koyanagi Y, Mitsuya H, Matsuoka M. (2006) De novo human T-cell leukemia virus type 1 infection of human lymphocytes in NOD-SCID, common gamma-chain knockout mice. *J Virol*, 80(21):10683-91.
33. Nakamura T, Miyakawa Y, Miyamura A, Yamane A, Suzuki H, Ito M, Ohnishi Y, Ishiwata N, Ikeda Y, Tsuruzoe N. (2006) A novel nonpeptidyl human c-Mpl activator stimulates human megakaryopoiesis and thrombopoiesis. *Blood*, 107(11):4300-4307.
34. Ninomiya M, Abe A, Yokozawa T, Ozeki K, Yamamoto K, Ito M, Ito M, Kiyoi H, Emi N, Naoe T. (2006) Establishment of a myeloid leukemia cell line, TRL-O1, with MLL-ENL fusion gene. *Cancer Genet Cytogenet.*, 169:1-11.
35. Yahata T, Yumino S, Seng Y, Miyatake H, Uno T, Muguruma Y, Ito M, Miyoshi H, Kato S, Hotta T, Ando K. (2006) Clonal analysis of thymus-repopulating cells presents direct evidence for self-renewal division of human hematopoietic stem cells. *Blood*, 108:2446-2454.

TO ORDERUS: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM**The CIEA NOG mouse®**

36. Watanabe S, Terashima K, Ohta S, Horibata S, Yajima M, Shiozawa Y, Dewan MZ, Yu Z, Ito M, Morio T, Shimizu N, Honda M, Yamamoto N. (2007) Hematopoietic stem cell-engrafted NOD/SCID/IL2Rgamma null mice develop human lymphoid systems and induce long-lasting HIV-1 infection with specific humoral immune responses. *Blood*, 109:212-218.
37. Ninomiya M, Abe A, Katsumi A, Xu J, Ito M, Arai F, Suda T, Ito M, Kiyoi H, Kinoshita T, Naoe T. (2007) Homing, proliferation and survival sites of human leukemia cells *in vivo* in immunodeficient mice. *Leukemia*, 21:136-142.
38. Suemizu H, Monnai M, Ohnishi Y, Ito M, Tamaoki N, Nakamura M. (2007) Identification of a key molecular regulator of liver metastasis in human pancreatic carcinoma using a novel new quantitative model of metastasis in NOD/SCID/gammacnull (NOG) mice. *Int J Oncol.*, 31(4):741-51.
39. Watanabe M, Dewan MZ, Taira M, Shoda M, Honda M, Sata T, Higashihara M, Kadin ME, Watanabe T, Yamamoto N, Umezawa K, Horie R. (2007) I kappa B independent induction of NF-kappa B and its inhibition by DHMEQ in Hodgkin/Reed-Sternberg cells. *Lab Invest.*, 87:372-382.
40. Watanabe S, Ohta S, Yajima M, Terashima K, M.Ito, H.Mugishima H, Fujiwara S, Shimizu K, Honda M, Shimizu N, Yamamoto N (2007) Humanized NOD/SCID/IL2R{gamma}null mice transplanted with hematopoietic stem cells under non-myeloablative condition show prolonged life spans and allow detailed analysis of HIV-1 pathogenesis. *J Virol.*, 13259-13264.
41. Masuda H, Maruyama T, Hiratsu E, Yamane J, Iwanami A, Nagashima T, Ono M, Miyoshi H, Okano HJ, Ito M, Tamaoki N, Nomura T, Okano H, Matsuzaki Y, Yoshimura Y. (2007) Noninvasive and real-time assessment of reconstructed functional human endometrium in NOD/SCID/gammac(null) immunodeficient mice. *Proc Natl Acad Sci U S A.*, 104:1925-1930.
42. Ono, M, Maruyama T, Masuda H, Kajitani T, Nagashima T, Arase T, Ito M, Ohta K, Uchida H, Asada H, Yoshimura Y, Okano H, Matsuzaki Y. (2007) Side population in human uterine myometrium displays phenotypic and functional characteristics of myometrial stem cells. *Proc Natl Acad Sci U S A.*, 104(47):18700-5.
43. Fujino H, Hiramatsu H, Tsuchiya A, Niwa A, Noma H, Shiota M, Umeda K, Yoshimoto M, Ito M, Heike T, Nakahata T. (2007) Human cord blood CD34+ cells develop into hepatocytes in the livers of NOD/SCID/gamma(c)null mice through cell fusion. *FASEB J.*, 21:3499-3510.
44. Takajo I, Umeki K, Morishita K, Yamamoto I, Kubuki Y, Hatakeyama K, Kataoka H, Okayama A. (2007) Engraftment of peripheral blood mononuclear cells from human T-lymphotropic virus type 1 carriers in NOD/SCID/gammac(null) (NOG) mice. *Int J Cancer*, 121(10):2205-11.
45. Takenaka K, Prasolava TK, Wang JC, Mortin-Toth SM, Khalouei S, Gan OI, Dick JE, Danska JS. (2007) Polymorphism in Sirpa modulates engraftment of human hematopoietic stem cells. *Nat Immunol.* 8(12):1313-23.
46. Dewan MZ, Takamatsu N, Hidaka T, Hatakeyama K, Nakahata S, Fujisawa J, Katano H, Yamamoto N, Morishita K. (2008) Critical role for TSLC1 expression in the growth and organ infiltration of adult T-cell leukemia cells *in vivo*. *J Virol* 82:11958-11963.
47. Fujii E, Suzuki M, Matsubara K, Watanabe M, Chen YJ, Adachi K, Ohnishi Y, Tanigawa M, Tsuchiya M, Tamaoki N. (2008) Establishment and characterization of *in vivo* human tumor models in the NOD/SCID/gamma(c)(null) mouse. *Pathol Int* 58(9):559-67.
48. Hamada K, Monnai M, Kawai K, Nishime C, Kito C, Miyazaki N, Ohnishi Y, Nakamura M, Suemizu H. (2008) Liver metastasis models of colon cancer for evaluation of drug efficacy using NOD/Shi-scld IL2Rgammannull (NOG) mice. *J Oncol.*, 32:153-159.
49. Ito M, Kobayashi K, Nakahata T. (2008) NOD/Shi-scld IL2rgammac(null) (NOG) mice more appropriate for humanized mouse models. *Curr Top Microbiol Immunol*, 324:53-76.
50. Ito R, Shiina M, Saito Y, Tokuda Y, Kametani Y, Habu S. (2008) Antigen-specific antibody production of human B cells in NOG mice reconstituted with the human immune system. *Curr Top Microbiol Immunol*, 324:95-107.
51. Kishi Y, Tanaka Y, Shibata H, Nakamura S, Takeuchi K, Masuda S, Ikeda T, Muramatsu S, Hanazono Y. (2008) Variation in the incidence of teratomas after the transplantation of nonhuman primate ES cells into immunodeficient mice. *Cell Transplant*, 17(9):1095-102.
52. Koyanagi Y, Tanaka Y, Ito M, Yamamoto N. (2008) Humanized mice for human retrovirus infection. *Curr Top Microbiol Immunol*, 324:133-48.
53. Miyake A, Dewan MZ, Ishida T, Watanabe M, Honda M, Sata T, Yamamoto N, Umezawa K, Watanabe T, Horie R. (2008) Induction of apoptosis in Epstein-Barr virus-infected B-lymphocytes by the NF-kappa B inhibitor DHMEQ. *Microbes Infect* 10:748-756.

54. Nakamura M, Suemizu H. (2008) Novel metastasis models of human cancer in NOG mice. *Curr Top Microbiol Immunol*, 324:167-77.
55. Nogami W, Yoshida H, Koizumi K, Yamada H, Abe K, Arimura A, Yamane N, Takahashi K, Yamane A, Oda A, Tanaka Y, Takemoto H, Ohnishi Y, Ikeda Y, Miyakawa Y. (2008) The effect of a novel, small non-peptidyl molecule butyzamide on human thrombopoietin receptor and megakaryopoiesis. *Haematologica*, 93(10):1495-504.
56. Nomura T, Tamaoki N, Takakura A, Suemizu H. (2008) Basic concept of development and practical application of animal models for human diseases. *Curr Top Microbiol Immunol*, 324:1-24.
57. Suemizu H, Yagihashi C, Mizushima T, Ogura T, Etoh T, Kawai K, Ito M. (2008) Establishing EGFP congenic mice in a NOD/Shi-scld IL2Rg(null) (NOG) genetic background using a marker-assisted selection protocol (MASP). *Exp Anim*, 57(5):471-7.
58. Suemizu H, Hasegawa M, Kawai K, Taniguchi K, Monnai M, Wakui M, Suematsu M, Ito M, Peltz G, Nakamura M. (2008) Establishment of a humanized model of liver using NOD/Shi-scld IL2Rgnull mice. *Biochem Biophys Res Commun*, 377(1):248-52.
59. Terada Y, Terunuma-Sato Y, Kakoi-Yoshimoto T, Hasegawa H, Ugajin T, Koyanagi Y, Ito M, Murakami T, Sasano H, Yaegashi N, Okamura K. (2008) Development of human Graafian follicles following transplantation of human ovarian tissue into NOD/SCID/gammac null mice. *Am J Reprod Immunol*, 60:534-540.
60. Terunuma H, Deng X, Dewan Z, Fujimoto S, Yamamoto N. (2008) Potential role of NK cells in the induction of immune responses: implications for NK cell-based immunotherapy for cancers and viral infections. *Int Rev Immunol* 27:93-110.
61. Yahata T, Muguruma Y, Yumino S, Sheng Y, Uno T, Matsuzawa H, Ito M, Kato S, Hotta T, Ando K. (2008) Quiescent human hematopoietic stem cells in the bone marrow niches organize the hierarchical structure of hematopoiesis. *Stem Cells* 26:3228-3236.
62. Yajima M, Imadome K, Nakagawa A, Watanabe S, Terashima K, Nakamura H, Ito M, Shimizu N, Honda M, Yamamoto N, Fujiwara S. (2008) A new humanized mouse model of Epstein-Barr virus infection that reproduces persistent infection, lymphoproliferative disorder, and cell-mediated and humoral immune responses. *J Infect Dis*, 198(5):673-82.
63. Yamane A, Nakamura T, Suzuki H, Ito M, Ohnishi Y, Ikeda Y, Miyakawa Y. (2008) Interferon-alpha 2b-induced thrombocytopenia is caused by inhibition of platelet production but not proliferation and endomitosis in human megakaryocytes. *Blood*, 112(3):542-50.
64. Chijiwa T, Abe Y, Ikoma N, Yamazaki H, Tsukamoto H, Suemizu H, Kawai K, Wakui M, Nishime C, Matsumoto H, Matsuyama M, Mukai M, Ueyama Y, Nakamura M. (2009) Thrombospondin 2 inhibits metastasis of human malignant melanoma through microenvironment-modification in NOD/SCID/gammaCnull (NOG) mice. *Int J Oncol.*, 34(1):5-13.
65. Ito A, Ishida T, Yano H, Inagaki A, Suzuki S, Sato F, Takino H, Mori F, Ri M, Kusumoto S, Komatsu H, Iida S, Inagaki H, Ueda R. (2009) Defucosylated anti-CCR4 monoclonal antibody exercises potent ADCC-mediated antitumor effect in the novel tumor-bearing humanized NOD/Shi-scld, IL-2Rgamma(null) mouse model. *Cancer Immunol Immunother*, 58(8):1195-206.
66. Machida K, Suemizu H, Kawai K, Ishikawa T, Sawada R, Ohnishi Y, Tsuchiya T. (2009) Higher susceptibility of NOG mice to xenotransplanted tumors. *J Toxicol Sci*, 34:123-127.
67. Ogawa D, Okada Y, Nakamura M, Kanemura Y, Okano HJ, Matsuzaki Y, Shimazaki T, Ito M, Ikeda E, Tamiya T, Nagao S, Okano H. (2009) Evaluation of human fetal neural stem/progenitor cells as a source for cell replacement therapy for neurological disorders: properties and tumorigenicity after long-term *in vitro* maintenance. *J Neuro Res* 87:307-317.
68. Oh JY, Kim MK, Ko JH, Lee HJ, Kim Y, Park CS, Park CG, Kim SJ, Wee WR, Lee JH. (2009) Acute cell-mediated rejection in orthotopic pig-to-mouse corneal xenotransplantation. *Xenotransplantation.*, 16(2):74-82.
69. Ito R, Katano I, Kawai K, Hirata H, Ogura T, Kamisako T, Eto T, Ito M. (2009) Highly sensitive model for xenogenic GVHD using severe immunodeficient NOG mice. *Transplantation*, 87(11):1654-8.
70. Kato C, Fujii E, Chen YJ, Endaya BB, Matsubara K, Suzuki M, Ohnishi Y, Tamaoki N. (2009) Spontaneous thymic lymphomas in the non-obese diabetic/Shi-scld, IL-2R{gamma}null mouse. *Lab Anim*. 2009 Oct;43(4):402-4.
71. Inoue M, Senju S, Hirata S, Irie A, Baba H, Nishimura Y. (2009) An *in vivo* model of priming of antigen-specific human CTL by Mo-DC in NOD/Shi-scld IL2rgamma(null) (NOG) mice. *Immunol Lett*, 126(1-2):67-72.
72. Tanaka Y, Ikeda T, Kishi Y, Masuda S, Shibata H, Takeuchi K, Komura M, Iwanaka T, Muramatsu S, Kondo Y, Takahashi K, Yamanaka S, Hanazono Y. (2009) ERas is expressed in primate embryonic stem cells but not related to tumorigenesis. *Cell Transplant*, 18(4):381-9.

TO ORDERUS: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

73. Watanabe Y, Takahashi T, Okajima A, Shiokawa M, Ishii N, Katano I, Ito R, Ito M, Minegishi M, Minegishi N, Tsuchiya S, Sugamura K. (2009) The analysis of the functions of human B and T cells in humanized NOD/shi-scid/gammac(null) (NOG) mice (hu-HSC NOG mice). *Int Immunol*, 21(7):843-58.
74. Kawaguchi AT, Kametani Y, Kato S, Furuya H, Tamaoki K, Habu S. (2009) Effects of liposome-encapsulated hemoglobin on human immune system: evaluation in immunodeficient mice reconstituted with human cord blood stem cells. *Artif Organs*, 33(2):169-76.
75. Watanabe T, Hayashi H, Kita K, Kubota Y, Ogawa T. (2009) Ectopic porcine spermatogenesis in murine subcutis: tissue grafting versus cell-injection methods. *Asian J Androl*, 11(3):317-23.
76. Kai M, Hagiwara T, Emuta C, Chisaka Y, Tsuruhata K, Endo C, Inagaki Y, Miyazaki H, Kataoka S. (2009) *In vivo* efficacy of anti-MPL agonist antibody in promoting primary human hematopoietic cells. *Blood*, 113(10):2213-6.
77. Shima H, Takubo K, Iwasaki H, Yoshihara H, Gomei Y, Hosokawa K, Arai F, Takahashi T, Suda T. (2009) Reconstitution activity of hypoxic cultured human cord blood CD34-positive cells in NOG mice. *Biochem Biophys Res Commun*, 378(3):467-72.
78. Ishigaki T, Sudo K, Hiroyama T, Mihaara K, Ninomiya H, Chiba S, Nagasawa T, Nakamura Y. (2009) Human hematopoietic stem cells can survive *in vitro* for several months. *Adv Hematol* 2009: 936761.
79. Ito A, Ishida T, Utsunomiya A, Sato F, Mori F, Yano H, Inagaki A, Suzuki S, Takino H, Ri M, Kusumoto S, Komatsu H, Iida S, Inagaki H, Ueda R. (2009) Defucosylated anti-CCR4 monoclonal antibody exerts potent ADCC against primary ATLL cells mediated by autologous human immune cells in NOD/Shi-scid, IL-2R gamma(null) mice *in vivo*. *J Immunol* 183(7):4782-91.
80. Nie C, Sato K, Misawa N, Kitayama H, Fujino H, Hiramatsu H, Heike T, Nakahata T, Tanaka Y, Ito M, Koyanagi Y. (2009) Selective infection of CD4⁺ effector memory T lymphocytes leads to preferential depletion of memory T lymphocytes in R5 HIV-1-infected humanized NOD/SCID/IL-2Rgamma null mice. *Virology* 394:64-72.
81. Yajima M, Imadome K, Nakagawa A, Watanabe S, Terashima K, Nakamura H, Ito M, Shimizu N, Yamamoto N, Fujiwara S. (2009) T cell-mediated control of Epstein-Barr virus infection in humanized mice. *J Infect Dis* 200:1611-1615.
82. Ogawa D, Okada Y, Nakamura M, Kanemura Y, Okano HJ, Matsuzaki Y, Shimazaki T, Ito M, Ikeda E, Tamiya T, Nagao S, Okano H. (2009) Evaluation of human fetal neural stem/progenitor cells as a source for cell replacement therapy for neurological disorders: properties and tumorigenicity after long-term *in vitro* maintenance. *J Neurosci Res* 87:307-317.
83. Tanizaki R, Nomura Y, Miyata Y, Minami Y, Abe A, Hanamura A, Sawa M, Murata M, Kiyoi H, Matsushita T, Naoe T. (2009) Irrespective of CD34 expression, lineage-committed cell fraction reconstitutes and re-establishes transformed Philadelphia chromosome-positive leukemia in NOD/SCID/IL-2Rgamma^{-/-} mice. *Cancer Sci* 101(3):631-638.
84. Dewan MZ, Tomita M, Katano H, Yamamoto N, Ahmed S, Yamamoto M, Sata T, Mori N. (2009) An HIV protease inhibitor, ritonavir targets the nuclear factor-kappaB and inhibits the tumor growth and infiltration of EBV-positive lymphoblastoid B cells. *Int J Cancer* 124:622-629.
85. Handa K, Ohmura M, Nishime C, Hishiki T, Nagahata Y, Kawai K, Suemizu H, Nakamura M, Wakui M, Kitagawa Y, Suematsu M, Tsukada K. (2010) Phosphorescence-assisted microvascular O₂ measurements reveal alterations of oxygen demand in human metastatic colon cancer in the liver of superimmunodeficient NOG mice. *Adv Exp Med Biol* 662:423-9.
86. Kikuchi-Arai M, Murakami T, Utsunomiya H, Akahira J, Suzuki-Kakisaka H, Terada Y, Tachibana M, Hayasaka S, Ugajin T, Yaegashi N. (2010) Establishment of long-term model throughout regular menstrual cycles in immunodeficient mice. *Am J Reprod Immunol* 64:324-332.
87. Kamiyama H, Kamiyama M, Hong SM, Karikari CA, Lin MT, Borges MW, Griffith M, Young A, Norris- Kirby A, Lubek C, Mizuma M, Feldmann G, Shi C, Liang H, Goggins MG, Maitra A, Hruban RH, Eshleman JR. (2010) *In vivo* and *in vitro* propagation of intraductal papillary mucinous neoplasms. *Lab Invest* 90:665-673.
88. Kimura O, Takahashi T, Ishii N, Inoue Y, Ueno Y, Kogure T, Fukushima K, Shiina M, Yamagiwa Y, Kondo Y, Inoue J, Kakazu E, Iwasaki T, Kawagishi N, Shimosegawa T, Sugamura K. (2010) Characterization of the epithelial cell adhesion molecule (EpCAM)⁺ cell population in hepatocellular carcinoma cell lines. *Cancer Sci* 101:2145-2155.
89. Masuda H, Matsuzaki Y, Hiratsu E, Ono M, Nagashima T, Kajitani T, Arase T, Oda H, Uchida H, Asada H, Ito M, Yoshimura Y, Maruyama T, Okano H. (2010) Stem cell-like properties of the endometrial side population: implication in endometrial regeneration. *PLoS One* 5:e10387.
90. Tsuiji K, Takeda T, Li B, Kondo A, Ito M, Yaegashi N. (2010) Establishment of a novel xenograft model for human uterine leiomyoma in immunodeficient mice. *Tohoku J Exp Med* 222:55-61.

91. Sato K, Nie C, Misawa N, Tanaka Y, Ito M, Koyanagi Y. (2010) Dynamics of memory and naive CD8+ T lymphocytes in humanized NOD/SCID/IL-2R γ null mice infected with CCR5-tropic HIV-1. *Vaccine* 28 Suppl 2:B32-37.
92. Sato K, Izumi T, Misawa N, Kobayashi T, Yamashita Y, Ohmichi M, Ito M, Takaori-Kondo A, Koyanagi Y. (2010) Remarkable lethal G-to-A mutations in *vif*-proficient HIV-1 provirus by individual APOBEC3 proteins in humanized mice. *J Virol* 84:9546-9556.
93. Shiokawa M, Takahashi T, Murakami A, Kita S, Ito M, Sugamura K, Ishii N. (2010) *In vivo* assay of human NK-dependent ADCC using NOD/SCID/ γ null (NOG) mice. *Biochem Biophys Res Commun* 399:733-737.
94. Wakui M, Suemizu H. (2010) [Development of humanized models using NOG mice]. *Seikagaku* 82(4):314-8. [Article in Japanese]
95. Yamamoto I, Takajo I, Umeki K, Morishita K, Hatakeyama K, Kataoka H, Nomura H, Okayama A. (2010) Multiple integrations of human T-lymphotropic virus type 1 proviruses in the engrafted cells from the asymptomatic carriers in NOD/SCID/ γ null mice. *Intervirology* 53:229-239.
96. Zhang L, Meissner E, Chen J, Su L. (2010) Current humanized mouse models for studying human immunology and HIV-1 immuno-pathogenesis. *Sci China Life Sci* 53(2):195-203.
97. Asakura K, Izumi Y, Yamamoto M, Yamauchi Y, Kawai K, Serizawa A, Mizushima T, Ohmura M, Kawamura M, Wakui M, Adachi T, Nakamura M, Suematsu M, Nomori H. (2011) The cytostatic effects of lovastatin on ACC-MESO-1 cells. *J Surg Res* 170(2):e197-209.
98. Hasegawa M, Kawai K, Mitsui T, Taniguchi K, Monnai M, Wakui M, Ito M, Suematsu M, Peltz G, Nakamura M (2011) The reconstituted 'humanized liver' in TK-NOG mice is mature and functional. *Biochem Biophysical Res Communications* 405:405-410.
99. Hirose T, Torikai H, Yanagisawa M, Kamei M, Imahashi N, Demachi-Okamura A, Tanimoto M, Shiraiishi K, Ito M, Miyamura K, Shibata K, Kikkawa F, Morishima Y, Takahashi T, Emi N, Kuzushima K, Akatsuka Y. (2011) Mismatched human leukocyte antigen class II-restricted CD8(+) cytotoxic T cells may mediate selective graft-versus-leukemia effects following allogeneic hematopoietic cell transplantation. *Cancer Sci* 102(7):1281-6.
100. Katano I, Ito R, Eto T, Aiso S, Ito M. (2011) 14. Immunodeficient NOD-scid IL-2R γ null mice do not display T and B cell leakiness. *Exp Anim* 60(2):181-6.
101. Kubo A, Ohmura M, Wakui M, Harada T, Kajihara S, Ogawa K, Suemizu H, Nakamura M, Setou M, Suematsu M. (2011) Semi-quantitative analyses of metabolic systems of human colon cancer metastatic xenografts in livers of superimmunodeficient NOG mice. *Anal Bioanal Chem* 400(7):1895- 904.
102. Kuwana Y, Takei M, Yajima M, Imadome K, Inomata H, Shiozaki M, Ikumi N, Nozaki T, Shiraiwa H, Kitamura N, Takeuchi J, Sawada S, Yamamoto N, Shimizu N, Ito M, Fujiwara S. (2011) Epstein-Barr virus induces erosive arthritis in humanized mice. *PLoS One*. 6(10):e26630.
103. Muguruma Y, Matsushita H, Yahata T, Yumino S, Tanaka Y, Miyachi H, Ogawa Y, Kawada H, Ito M, Ando K. (2011) Establishment of a xenograft model of human myelodysplastic syndromes. *Haematologica* 96(4):543-51.
104. Sato K, Misawa N, Nie C, Satou Y, Iwakiri D, Matsuoka M, Takahashi R, Kuzushima K, Ito M, Takada K, Koyanagi Y. (2011) A novel animal model of Epstein-Barr virus-associated hemophagocytic lymphohistiocytosis in humanized mice. *Blood* 117(21):5663-73.
105. Yahata T, Takanashi T, Muguruma Y, Ibrahim AA, Matsuzawa H, Uno T, Sheng Y, Onizuka M, Ito M, Kato S, Ando K. (2011) Accumulation of oxidative DNA damage restricts the self-renewal capacity of human hematopoietic stem cells. *Blood* 118(11):2941-50.
106. Iriuchishima H, Takubo K, Miyakawa Y, Nakamura-Ishizu A, Miyauchi Y, Fujita N, Miyamoto K, Miyamoto T, Ikeda E, Kizaki M, Nojima Y, Suda T. (2012) Neovascular niche for human myeloma cells in immunodeficient mouse bone. *PLoS One*. 7(2):e30557.
107. Ito R, Katano I, Ida-Tanaka M, Kamisako T, Kawai K, Suemizu H, Aiso S, Ito M. (2012) Efficient xenoengraftment in severe immunodeficient NOD/Shi-*scid* IL2r γ null mice is attributed to a lack of CD11c⁺B220⁺CD122⁺ cells. *J Immunol*. 189(9):4313-20.

TO ORDERUS: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

108. Kobayashi S, Yamada-Okabe H, Suzuki M, Natori O, Kato A, Matsubara K, Jau Chen Y, Yamazaki M, Funahashi S, Yoshida K, Hashimoto E, Watanabe Y, Mutoh H, Ashihara M, Kato C, Watanabe T, Yoshikubo T, Tamaoki N, Ochiya T, Kuroda M, Levine AJ, Yamazaki T. (2012) LGR5-positive colon cancer stem cells interconvert with drug-resistant LGR5-negative cells and are capable of tumor reconstitution. *Stem Cells*. 30(12):2631-44.
109. Kondo S, Iwata S, Yamada T, Inoue Y, Ichihara H, Kichikawa Y, Katayose T, Souta-Kuribara A, Yamazaki H, Hosono O, Kawasaki H, Tanaka H, Hayashi Y, Sakamoto M, Kamiya K, Dang NH, Morimoto C. (2012) Impact of the integrin signaling adaptor protein NEDD9 on prognosis and metastatic behavior of human lung cancer. *Clin Cancer Res*. 18(22):6326-38.
110. Mori F, Ishida T, Ito A, Sato F, Masaki A, Takino H, Ri M, Kusumoto S, Komatsu H, Ueda R, Inagaki H, Iida S. (2012) Potent antitumor effects of bevacizumab in a microenvironment-dependent human lymphoma mouse model. *Blood Cancer J*. 2(4):e67.
111. Moriya K, Suzuki M, Watanabe Y, Takahashi T, Aoki Y, Uchiyama T, Kumaki S, Sasahara Y, Minegishi M, Kure S, Tsuchiya S, Sugamura K, Ishii N. (2012) Development of a multi-step leukemogenesis model of MLL-rearranged leukemia using humanized mice. *PLoS One*. 7(6):e37892.
112. Nischang M, Suttmuller R, Gers-Huber G, Audigé A, Li D, Rochat MA, Baenziger S, Hofer U, Schlaepfer E, Regenass S, Amssoms K, Stoops B, Van Cauwenberge A, Boden D, Kraus G, Speck RF. (2012) Humanized mice recapitulate key features of HIV-1 infection: a novel concept using long-acting anti-retroviral drugs for treating HIV-1. *PLoS One*. 7(6):e38853.
113. Nunomura S, Shimada S, Kametani Y, Yamada Y, Yoshioka M, Suemizu H, Ozawa M, Itoh T, Kono A, Suzuki R, Tani K, Ando K, Yagita H, Ra C, Habu S, Satake M, Sasaki E. (2012) Double expression of CD34 and CD117 on bone marrow progenitors is a hallmark of the development of functional mast cell of *Callithrix jacchus* (common marmoset). *Int Immunol*. 24(9):593-603.
114. Saito N, Yoshioka N, Abe R, Qiao H, Fujita Y, Hoshina D, Suto A, Kase S, Kitaichi N, Ozaki M, Shimizu H. (2012) Stevens-Johnson syndrome/toxic epidermal necrolysis mouse model generated by using PBMCs and the skin of patients. *J Allergy Clin Immunol*. 131(2):434-41.e1-9.
115. Schoenbrunn A, Frentsch M, Kohler S, Keye J, Dooms H, Moewes B, Dong J, Loddenkemper C, Sieper J, Wu P, Romagnani C, Matzmohr N, Thiel A. (2012) A converse 4-1BB and CD40 ligand expression pattern delineates activated regulatory T cells (Treg) and conventional T cells enabling direct isolation of alloantigen-reactive natural Foxp3+ Treg. *J Immunol*. 189(12):5985-94.
116. Shirakura Y, Mizuno Y, Wang L, Imai N, Amaike C, Sato E, Ito M, Nukaya I, Mineno J, Takesako K, Ikeda H, Shiku H. (2012) T-cell receptor gene therapy targeting melanoma-associated antigen-A4 inhibits human tumor growth in non-obese diabetic/SCID/γcnull mice. *Cancer Sci*. 103(1):17-25.
117. Takahashi M, Tsujimura N, Otsuka K, Yoshino T, Mori T, Matsunaga T, Nakasono S. (2012) Comprehensive evaluation of leukocyte lineage derived from human hematopoietic cells in humanized mice. *J Biosci Bioeng*. 113(4):529-35.
118. Takahashi M, Tsujimura N, Yoshino T, Hosokawa M, Otsuka K, Matsunaga T, Nakasono S. (2012) Assessment of benzene-induced hematotoxicity using a human-like hematopoietic lineage in NOD/Shi-scid/IL-2Rγnull mice. *PLoS One*. 7(12):e50448.
119. Zhang Y, Patel S, Abdelouahab H, Wittner M, Willekens C, Shen S, Betems A, Joulin V, Opolon P, Bawa O, Pasquier F, Ito M, Fujii N, Gonin P, Solary E, Vainchenker W, Coppo P, De Botton S, Louache F. (2012) CXCR4 inhibitors selectively eliminate CXCR4-expressing human acute myeloid leukemia cells in NOG mouse model. *Cell Death Dis*. 3:e396.
120. Fukuda K, Saikawa Y, Sako H, Yoshimura Y, Takahashi T, Wada N, Kawakubo H, Takeuchi H, Ohmori T, Kitagawa Y. (2013) Establishment and characterization of novel cell lines and xenografts from patients with gastrointestinal stromal tumors. *Oncol Rep*. 30(1):71-8.
121. Joo KM, Kim J, Jin J, Kim M, Seol HJ, Muradov J, Yang H, Choi YL, Park WY, Kong DS, Lee JI, Ko YH, Woo HG, Lee J, Kim S, Nam DH. (2013) Patient-specific orthotopic glioblastoma xenograft models recapitulate the histopathology and biology of human glioblastomas in situ. *Cell Rep*. 3(1):260-73.
122. Kanemura H, Go MJ, Nishishita N, Sakai N, Kamao H, Sato Y, Takahashi M, Kawamata S. (2013) Pigment Epithelium-Derived Factor Secreted from Retinal Pigment Epithelium Facilitates Apoptotic Cell Death of iPSC. *Sci Rep*. 3:2334.
123. Kato T, Hisasue M, Segawa K, Fujimoto A, Makiishi E, Neo S, Yasuno K, Kobayashi R, Tsuchiya R. (2013) Accumulation of Xenotransplanted Canine Bone Marrow Cells in NOD/SCID/γc(null) Mice with Acute Hepatitis Induced by CCl4. *J Vet Med Sci*. 75(7):847-55.

124. Kawai K, Tamura K, Sakata I, Ishida J, Nagata M, Tsukada H, Suemizu H, Nakamura M, Abe Y, Chijiwa T. (2013) A new *in vivo* model to analyze hepatic metastasis of the human colon cancer cell line HCT116 in NOD/Shi-scid/IL-2R γ (null) (NOG) mice by (18)F-FDG PET/CT. *Oncol Rep.* 29(2):464-8.
125. Masaki A, Ishida T, Suzuki S, Ito A, Mori F, Sato F, Narita T, Yamada T, Ri M, Kusumoto S, Komatsu H, Tanaka Y, Niimi A, Inagaki H, Iida S, Ueda R. (2013) Autologous Tax-specific CTL therapy in a primary adult T cell leukemia/lymphoma cell-bearing NOD/Shi-scid, IL-2R γ null mouse model. *J Immunol.* 191(1):135-44.
126. Matsuzaki J, Suzuki H, Tsugawa H, Watanabe M, Hossain S, Arai E, Saito Y, Sekine S, Akaike T, Kanai Y, Mukaisho KI, Auwerx J, Hibi T. (2013) Bile Acids Increase Levels of microRNAs 221 and 222, Leading to Degradation of CDX2 During Esophageal Carcinogenesis. *Gastroenterology.* S0016- 5085(13)01154-2.
127. Murata T, Iwata S, Siddiquey MN, Kanazawa T, Goshima F, Kawashima D, Kimura H, Tsurumi T. (2013) Heat shock protein 90 inhibitors repress latent membrane protein 1 (LMP1) expression and proliferation of Epstein-Barr virus-positive natural killer cell lymphoma. *PLoS One.* 8(5):e63566.
128. Reghunathan R, Bi C, Liu SC, Loong KT, Chung TH, Huang G, Chng WJ. (2013) Clonogenic multiple myeloma cells have shared stemness signature associated with patient survival. *Oncotarget.* 4(8):1230-40.
129. Shimizu K, Mizuno T, Shinga J, Asakura M, Kakimi K, Ishii Y, Masuda K, Maeda T, Sugahara H, Sato Y, Matsushita H, Nishida K, Hanada K, Dorrie J, Schaft N, Bickham K, Koike H, Ando T, Nagai R, Fujii S. (2013) Vaccination with antigen-transfected, NKT cell ligand-loaded, human cells elicits robust *in situ* immune responses by dendritic cells. *Cancer Res.* 73(1):62-73.
130. Wang C, Nakamura S, Oshima M, Mochizuki-Kashio M, Nakajima-Takagi Y, Osawa M, Kusunoki Y, Kyoizumi S, Imai K, Nakachi K, Iwama A. (2013) Compromised hematopoiesis and increased DNA damage following non-lethal ionizing radiation of a human hematopoietic system reconstituted in immunodeficient mice. *Int J Radiat Biol.* 89(2):132-7.
131. Yoshimura H, Matsuda Y, Naito Z, Ishiwata T (2013) *In vivo* bioluminescence imaging of pancreatic cancer xenografts in NOG mice. *J Carcinogene Mutagene* S9:003.
132. Nakatsuka R, Matsuoka Y, Uemura Y, Sumide K, Iwaki R, Takahashi M, Fujioka T, Sasaki Y, Sonoda Y. (2013) Mouse dental pulp stem cells support human umbilical cord blood-derived hematopoietic stem/progenitor cells *in vitro*. *Cell Transplant.* Oct 29.
133. Søndergaard H, Kvist PH, Haase C. (2013) Human T cells depend on functional calcineurin, tumour necrosis factor- α and CD80/CD86 for expansion and activation in mice. *Clin Exp Immunol.* 172(2):300-10.
134. Tomita H, Fuchimoto Y, Mori T, Kato J, Uemura T, Handa M, Tazawa H, Ohdan H, Okamoto S, Kuroda T. (2013) Production of anti-ABO blood group antibodies after minor ABO-incompatible bone marrow transplantation in NOD/SCID/gamma(c)(null) mice. *Clin Transplant.* 27(6):E702-8.
135. Imadome K. (2013) A novel mouse xenotransplantation model of EBV-T/NK-LPD and the application of the mouse model. *Nihon Rinsho Meneki Gakkai Kaishi.* 36(6):433-41.
136. Mori F, Ishida T, Ito A, Sato F, Masaki A, Narita T, Suzuki S, Yamada T, Takino H, Ri M, Kusumoto S, Komatsu H, Hishizawa M, Imada K, Takaori-Kondo A, Niimi A, Ueda R, Inagaki H, Iida S. (2014) Antitumor effects of bevacizumab in a microenvironment-dependent human adult T-cell leukemia/lymphoma mouse model. *Eur J Haematol.* 92(3):219-28.
137. Hertlein E, Beckwith KA, Lozanski G, Chen TL, Towns WH, Johnson AJ, Lehman A, Ruppert AS, Bolon B, Andritsos L, Lozanski A, Rassenti L, Zhao W, Jarvinen TM, Senter L, Croce CM, Symer DE, de la Chapelle A, Heerema NA, Byrd JC. (2013) Characterization of a new chronic lymphocytic leukemia cell line for mechanistic *in vitro* and *in vivo* studies relevant to disease. *PLoS One.* 8(10):e76607.
138. Tezuka K, Xun R, Tei M, Ueno T, Tanaka M, Takenouchi N, Fujisawa J. (2014) An animal model of adult T-cell leukemia: humanized mice with HTLV-1-specific immunity. *Blood.* 123(3):346-55.
139. Matsuda Y, Yoshimura H, Ueda J, Naito Z, Korc M, Ishiwata T. (2014) Nestin delineates pancreatic cancer stem cells in metastatic foci of NOD/Shi-scid IL2R γ (null) (NOG) mice. *Am J Pathol.* 184(3):674-85.
140. Fujii E, Kato A, Chen YJ, Matsubara K, Ohnishi Y, Suzuki M. (2014) Histopathological characteristics of human non-tumor thyroid tissues in a long-term model of adenomatous goiter xenografts in the NOD/Shi-scid, IL-2R γ (null) mouse. *Exp Toxicol Pathol.* 66(4):203-9.

TO ORDER

US: 1-888-822-6642 | EU: +45 70 23 04 05 | INFO@TACONIC.COM

The CIEA NOG mouse®

141. Borriello L, Montès M, Lepelletier Y, Leforban B, Liu WQ, Demange L, Delhomme B, Pavoni S, Jarray R, Boucher JL, Dufour S, Hermine O, Garbay C, Hadj-Slimane R, Raynaud F. (2014) Structure-based discovery of a small non-peptidic Neuropilins antagonist exerting *in vitro* and *in vivo* anti-tumor activity on breast cancer model. *Cancer Lett.* 349(2):120-7.
142. Haji Y, Suzuki M, Moriya K, So T, Hozumi K, Mizuma M, Unno M, Ishii N. (2014) Activation of Notch1 promotes development of human CD8(+) single positive T cells in humanized mice. *Biochem Biophys Res Commun.* 447(2):346-51.
143. Tamai K, Nakamura M, Mizuma M, Mochizuki M, Yokoyama M, Endo H, Yamaguchi K, Nakagawa T, Shiina M, Unno M, Muramoto K, Sato I, Satoh K, Sugamura K, Tanaka N. (2014) Suppressive expression of CD274 increases tumorigenesis and cancer stem cell phenotypes in cholangiocarcinoma. *Cancer Sci.*;105(6):667-74.
144. Poillet L, Pernodet N, Boyer-Guittaut M, Adami P, Borg C, Jouvenot M, Delage-Mourroux R, Despouy G. (2014) QSOX1 inhibits autophagic flux in breast cancer cells. *PLoS One.* 9(1):e86641.

To view an updated listing of references describing the use of the NOG mouse, organized by application visit

[Taconic.com/resources/nog-publications](https://www.taconic.com/resources/nog-publications)

Take Your Research Further



GEMS DESIGN

Taconic Biosciences GEMs Design empowers our clients to develop research models specifically suited to the unique needs of their discovery and development studies or therapeutic programs.

- Gene Inactivation
- Gene Mutation or Replacement
- CRISPR Gene Editing
- Transgene Expression
- miRNA Expression
- Cohort Production Packages

PRECISION RESEARCH MODELS

Research organizations demand precision tools that better reflect human physiology. Taconic Biosciences leads the field delivering innovative solutions to meet these continually evolving needs. Our core competencies include the delivery of complex strategies that both integrate human genetic sequences and engraft human cells and tissues into custom mouse and rat models.

- Human Gene Replacement
- Human Cell and Tissue Engraftment

GEMS MANAGEMENT

Taconic's fully integrated GEMs Management brings innovative models from design to study-ready cohorts with unprecedented speed and transparency.

- Embryology
- Rapid Colony Expansion
- Contract Breeding
- Surgical Services
- Tissue Collection
- Genotyping and Molecular Analysis
- Microbiome and Germ-Free Research Models and Services

CHOOSE TACONIC

For more than 60 years, Taconic has anticipated the needs of the scientific community to deliver models and services that meet the diverse needs of biomedical and biopharmaceutical researchers.

Today that forward thinking and commitment to working collaboratively has resulted in a client-centric environment infused with a knowledge bank that allows you to select the optimum model for your study based on informed insight into the generation of genetically engineered mouse and rat models.

YOUR COLLABORATIVE PARTNER

As a full-service biosciences company, Taconic can help you acquire, test, develop, breed, cryopreserve, prepare, and distribute highly relevant research lines worldwide. Whether you require custom genetically engineered, cell or tissue engrafted models or traditional models, Taconic's scientists will partner with you to rapidly and efficiently deliver the highest quality models.

TALK TO A SCIENTIST

Our scientific teams are happy to meet and talk with you about the most efficient way to achieve your study goals. Working in partnership with clients the world over, our scientific teams offer expert advice that can help you speed up your research and reduce your overall costs.

TALK TO A REPRESENTATIVE

For general information, you can talk to a member of our customer service team. Our customer service team is here to help you make the right decisions and get the models you need fast. Contact us at info@taconic.com

VISIT TACONIC.COM

For more information on the entire Taconic portfolio of products and services designed to help further your research, visit Taconic.com

©Taconic Biosciences, Inc. All rights reserved. Contents of this publication may not be reproduced in any form without prior permission.

US: 1-888-822-6642
EU: +45 70 23 04 05
INFO@TACONIC.COM

TACONIC.COM

0516

